

Superconformal Indices for $\mathcal{N} = 6$ Chern Simons Theories

Jyotirmoy Bhattacharya and Shiraz Minwalla

Department of Theoretical Physics, Tata Institute of Fundamental Research,
Homi Bhabha Rd, Mumbai 400005, India.

E-mail: jjyotirmoy@theory.tifr.res.in, minwalla@theory.tifr.res.in

ABSTRACT: Aharony, Bergman, Jafferis and Maldacena have recently proposed a dual gravitational description for a family of superconformal Chern Simons theories in three spacetime dimensions. In this note we perform the one loop computation that determines the field theory superconformal index of this theory and compare with the index computed over the Fock space of dual supersymmetric gravitons. In the appropriate limit (large N and large k) we find a perfect match.

Contents

1. Introduction	1
2. Field Theory computation of the index	3
3. Gravity computation of the index	7
4. Discussion	10

1. Introduction

Aharony, Bergman, Jaffiers and Maldacena (ABJM) have recently proposed that a class of $d = 3$ $U(N) \times U(N)$ $\mathcal{N} = 6$ superconformal Chern Simons field theories admit a dual description in terms of M theory compactified on $AdS_4 \times \frac{S^7}{Z_k}$ [1]. This correspondence has been further studied in [2]. The theories studied by these authors are parameterized by two integers N and k . In dual bulk terms k is the rank of the orbifold action on S^7 while N represents the number of units of 7 form flux that pierce S^7/Z_k . In field theory terms N denotes the rank of each of the $U(N)$ factors of the gauge group and $\pm k$ are the levels of the Chern Simons terms associated with each of these gauge groups.

With an appropriate normalization for fields, the effective 't Hooft coupling constant of any large N gauge theory is given by N times the inverse of the coefficient of the action. As the coefficient of the Chern Simons term is proportional to k , the effective 't Hooft coupling of the ABJM field theory is proportional to N/k . As N and k are both integers this 't Hooft coupling cannot be varied continuously; indeed a shift in k by unity shifts λ by the discrete amount $\delta\lambda = -\frac{\lambda^2}{N}$. Note however that $\delta\lambda \rightarrow 0$ in the 't Hooft limit ($N \rightarrow \infty$ with λ held fixed). Consequently λ is effectively a continuous parameter in the 't Hooft limit. It follows that the superconformal Witten index, defined for arbitrary 3 dimensional superconformal field theories in [3], must be invariant under deformations of λ in this 't Hooft scaling regime¹.

In this note we compute the superconformal index (as defined in [3]) of the ABJM theory in two different regimes. We first use the techniques of [4, 5, 6, 7, 8, 9] to find an expression for this index at $k = \infty$ (and so $\lambda = 0$ but at arbitrary N) in terms of

¹This is only true of contributions to the index from states whose energy stays finite as N is taken to infinity.

an integral over two $N \times N$ unitary matrices. Taking a further large N limit we are able to evaluate these unitary integrals explicitly using saddle point techniques.

Next we evaluate the index of this theory at infinite N and large λ , using the ABJM proposal for the dual description of this theory. Effectively, we compute the index over the Fock space of non interacting supersymmetric $U(1)$ neutral (see below) gravitons in $AdS_4 \times \frac{S^7}{Z_k}$. We perform this calculation explicitly in the 't Hooft limit, but also explain the generalization of this calculation to finite values of k , and so to values of λ that scale like N in the large N limit.

We find that our two independent computations of the superconformal Witten index of [3] agree perfectly in the 't Hooft limit. We view this agreement as a test of the ABJM proposal. Indeed, since the index computed in this paper is the most general superconformal index [3] for a $\mathcal{N} = 6$ superconformal field theory ² our calculation verifies the most detailed matching of supersymmetric states predicted by ABJM conjecture taken together with the requirement of superconformal invariance alone. Of course the agreement of the two independent index computations reported in this paper is closely related to the agreement of the spectrum of chiral operators of the field theory and the spectrum of gravitons in $AdS_4 \times \frac{S^7}{Z_k}$ reported in [1]. Note, however, that while the chiral ring described by ABJM counts states described by at least four supercharges (two supersymmetries and their Hermitian conjugates) the index constructed in this paper receives contributions from all states that are annihilated by a minimum of two supercharges. Note also that the invariance under λ deformations of the index computed in this paper follows from superconformal invariance alone, and uses no dynamical information about the field theory beyond its field content. In particular our field theoretic computation of the index is blind to the nature of the ABJM superpotential whose form played an important role in the construction of the ABJM chiral ring [1].

The matching of the index computed perturbatively in the ABJM theory with the index computed over the spectrum of bulk gravitons is reminiscent of a similar match in the case of $\mathcal{N} = 4$ Yang Mills theory [7]. As explained in that paper, $AdS_5 \times S^5$ hosts a family of 1/16 BPS black holes which the 4 dimensional superconformal index appears to be blind to. In a similar fashion the 3 dimensional superconformal index computed in this paper appears to be blind to the 1/12 BPS black holes presumably hosted by the $AdS_4 \times \frac{S^7}{Z_k}$ dual background (see [10, 11, 12]).

This note is organized as follows. In §2 below we review the symmetry algebra of the ABJM theory, the definition of the Witten index of [3] for this theory and the field content of the ABJM Chern Simons theory. We then perform a one loop field theory computation to present a field theoretic formula for this index in terms of an integral over two unitary matrices and evaluate these integrals in the large N limit. In §3 we present our computation of the same index over the spectrum of $U(1)$

²By a superconformal index we mean a quantity whose invariance under marginal deformations is guaranteed by superconformal invariance alone.

invariant multi gravitons in $AdS_4 \times \frac{S^7}{Z_k}$. In §4 we end with a discussion of our results and the generalizations they suggest.

2. Field Theory computation of the index

The symmetry algebra of the ABJM theory is the $d = 3$ $\mathcal{N} = 6$ superconformal algebra. The structure of this algebra and its unitary representations has been reviewed in detail in [3], and we will use the notation of that paper in what follows.

The bosonic subgroup of the $d = 3$ $\mathcal{N} = 6$ superconformal algebra is $SO(3, 2) \times SO(6)$. All states and operators in this theory are labeled by their quantum numbers under the maximal compact subalgebra of the superconformal algebra, $SO(3) \times SO(2) \times SO(6)$. In what follows we will denote the eigenvalue of the Cartan generator of $SO(3)$ by j , the eigenvalue under $SO(2)$ (the scaling dimension or global AdS_4 energy) by ϵ_0 and the three Cartan generators of $SO(6)$ - defined as the eigenvalues under the generators of rotations in the three orthogonal two planes - as h_1, h_2, h_3 .³

The twelve supercharges of the ABJM theory each have $\epsilon_0 = \frac{1}{2}$, and transform in the $j = \frac{1}{2}$ representation of $SO(3)$ algebra and the vector $(h_1, h_2, h_3) = (1, 0, 0)$ of $SO(6)$ algebra. The only propagating fields in the ABJM theory are a set of bi-fundamental and anti bi-fundamental scalars and fermions. All scalars have dimension $\epsilon_0 = \frac{1}{2}$ and are scalars under $SO(3)$, while all fermions have dimension $\epsilon_0 = 1$ and transform in the spin half representation of $SO(3)$. Bi-fundamental scalars/fermions transform in the $(\frac{1}{2}, \frac{1}{2}, \pm\frac{1}{2})$ representation of $SO(6)$, while anti bi-fundamental scalars and fermions transform in the $(\frac{1}{2}, \frac{1}{2}, \mp\frac{1}{2})$ representation of $SO(6)$. The symmetry transformation properties of the supersymmetries, propagating fields and derivatives of the ABJM theory are listed in table 1 below. In table 1 and throughout this paper, the symbol ϕ_{12} and ψ_{12} respectively denote scalar and fermionic fields that transform in the fundamental of the first $U(N)$ gauge group and antifundamental of the second $U(N)$ gauge group, while ϕ_{21} and ψ_{21} respectively denote scalar and fermionic fields which transforms in the antifundamental of the first $U(N)$ gauge group and in the fundamental of the second.

In this note we will compute the Witten index

$$I^W = \text{Tr} \left((-1)^F x^{\epsilon_0+j} y_1^{h_2} y_2^{h_3} \right) \quad (2.1)$$

for the ABJM theory quantized on $S^2 \times R$. As explained in [3] this index receives contributions only from states that are annihilated by a special supercharge Q together with its Hermitian conjugate $Q^\dagger$. Q has quantum numbers $\epsilon_0 = \frac{1}{2}$, $j = -\frac{1}{2}$,

³ $SO(6)$ may be thought of as the group of rotations about the origin in R^6 parameterized by x^i , $i = 1 \dots 6$. h_1 , h_2 and h_3 are simply the generators of rotations in the two planes (12), (34) and (56) respectively. Throughout this paper we will label representations of $SO(6)$ by their highest weights under (h_1, h_2, h_3) . In our conventions an $SO(6)$ weight is positive if h_1 is positive, or if $h_1 = 0$ and h_2 is positive or if $h_1 = h_2 = 0$ and h_3 is positive. We use a similar conventions for $SO(8)$ representations below.

Table 1: A list of the field content of ABJM theory, the supercharges, the derivatives and the representations.

type of operators	operators	scaling dimension (ϵ_0)	$SO(3)$ highest weight	$SO(6)$ highest weight
dynamical fields	ϕ_{12}	$\frac{1}{2}$	0	$(\frac{1}{2}, \frac{1}{2}, -\frac{1}{2})$
	ψ_{12}	1	$\frac{1}{2}$	$(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
	ϕ_{21}	$\frac{1}{2}$	0	$(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$
	ψ_{21}	1	$\frac{1}{2}$	$(\frac{1}{2}, \frac{1}{2}, -\frac{1}{2})$
supersymmetry generators	Q	$\frac{1}{2}$	$\frac{1}{2}$	(1, 0, 0)
derivatives	∂	1	1	(0, 0, 0)

$(h_1, h_2, h_3) = (1, 0, 0)$. It follows from the superconformal algebra that

$$\{Q, Q^\dagger\} = (\epsilon_0 - j - h_1) \equiv \Delta.$$

As a consequence, a state is annihilated by both Q and $Q^\dagger$ if and only if $\Delta = 0$. Consequently the index (2.1) receives contributions only from states with $\Delta = 0$. In table 2 below we list all $\Delta = 0$ propagating fields and derivatives of the ABJM theory. We also list the partition function over all $\Delta = 0$ bosonic fields and their derivatives (f_{12}^{bosonic} and f_{21}^{bosonic}), the partition function over $\Delta = 0$ fermionic fields and their derivatives ($f_{12}^{\text{fermionic}}$ and $f_{21}^{\text{fermionic}}$), and the Witten index (f_{12} and f_{21}) over these fields.

Following [4, 5] it was demonstrated in [8] (see equation(2.7)) that the free superconformal index of a Yang-Mills theory with the field content listed above is given by,

$$I^W = \int dU_1 dU_2 \exp \left(\sum_{n=1}^{\infty} \frac{1}{n} [f_{12}(x^n, y_1^n, y_2^n) \text{Tr} U_1^n \text{Tr} U_2^{-n} + f_{21}(x^n, y_1^n, y_2^n) \text{Tr} U_1^{-n} \text{Tr} U_2^n] \right) \quad (2.2)$$

The unitary integrals described in (2.2) may be evaluated in the large N limit. Let $\rho_n = \frac{\text{Tr} U_1^n}{N}$ and $\chi_n = \frac{\text{Tr} U_2^n}{N}$. In the large N limit the various ρ_n may be treated

Table 2: List of the supersymmetric ($\Delta = 0$) fields or ‘letters’ of the theory over which we calculate the index

letter	ϵ_0	$SO(3)$ (j)	$SO(6)$ weights	ϵ_0+j	partition function	index
bi-fundamental						
$(\phi_{12})_1$	$\frac{1}{2}$	0	$(\frac{1}{2}, \frac{1}{2}, -\frac{1}{2})$	$\frac{1}{2}$	$f_{12}^{\text{bosonic}} = \frac{x^{\frac{1}{2}}}{1-x^2} \left(\frac{y_1}{y_2} + \frac{y_1}{y_2} \right)$	$f_{12} = \frac{x^{\frac{1}{2}}}{1-x^2} \left(\frac{y_1}{y_2} + \frac{y_1}{y_2} \right) - \frac{x^{\frac{3}{2}}}{1-x^2} \left(y_1 y_2 + \frac{1}{y_1 y_2} \right)$
$(\phi_{12})_2$	$\frac{1}{2}$	0	$(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$	$\frac{1}{2}$		
$(\psi_{12})_1$	1	$\frac{1}{2}$	$(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$	$\frac{3}{2}$	$f_{12}^{\text{fermionic}} = \frac{x^{\frac{3}{2}}}{1-x^2} \left(y_1 y_2 + \frac{1}{y_1 y_2} \right)$	
$(\psi_{12})_2$	1	$\frac{1}{2}$	$(\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2})$	$\frac{3}{2}$		
anti bi-fundamental						
$(\phi_{21})_1$	$\frac{1}{2}$	0	$(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$	$\frac{1}{2}$	$f_{21}^{\text{bosonic}} = \frac{x^{\frac{1}{2}}}{1-x^2} \left(y_1 y_2 + \frac{1}{y_1 y_2} \right)$	$f_{21} = \frac{x^{\frac{1}{2}}}{1-x^2} \left(y_1 y_2 + \frac{1}{y_1 y_2} \right) - \frac{x^{\frac{3}{2}}}{1-x^2} \left(\frac{y_1}{y_2} + \frac{y_1}{y_2} \right)$
$(\phi_{21})_2$	$\frac{1}{2}$	0	$(\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2})$	$\frac{1}{2}$		
$(\psi_{21})_1$	1	$\frac{1}{2}$	$(\frac{1}{2}, \frac{1}{2}, -\frac{1}{2})$	$\frac{3}{2}$	$f_{21}^{\text{fermionic}} = \frac{x^{\frac{3}{2}}}{1-x^2} \left(\frac{y_1}{y_2} + \frac{y_1}{y_2} \right)$	
$(\psi_{21})_2$	1	$\frac{1}{2}$	$(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2})$	$\frac{3}{2}$		
derivative						
∂	1	+1	(0, 0, 0)	2		

as independent variables (modulo a positivity constraint - see for instance [5] - that

will turn out to be irrelevant for our considerations below) and

$$DU_1 = \prod_n d\rho_n \exp \left(-N^2 \sum_n \frac{\rho_n \rho_{-n}}{n} \right), \quad DU_2 = \prod_n d\rho_n \exp \left(-N^2 \sum_n \frac{\chi_n \chi_{-n}}{n} \right)$$

so that

$$I^W = \int \prod_{n \neq 0} d\rho_n d\chi_n \exp \left(N^2 \sum_{n=1}^{\infty} \frac{1}{n} (-|\rho_n|^2 - |\chi_n|^2 + f_{12}(x^n, y_1^n, y_2^n) \rho_n \chi_{-n} + f_{21}(x^n, y_1^n, y_2^n) \rho_{-n} \chi_n) \right). \quad (2.3)$$

The integral in (2.3) takes the form

$$I^W = \int \prod_{n \neq 0} d\rho_n d\chi_n \exp \left(-N^2 \sum_{n=1}^{\infty} \frac{1}{n} ((C^n)_i (M^n)_{ij} (C^n)_j) \right), \quad (2.4)$$

where the column C^n and the matrix $(M^n)_{ij}$ are given by

$$C^n = \begin{pmatrix} \chi_n \\ \rho_n \\ \chi_{-n} \\ \rho_{-n} \end{pmatrix}, \quad M^n = \frac{1}{2} \begin{pmatrix} 0 & 0 & 1 & -f_{21} \\ 0 & 0 & -f_{12} & 1 \\ 1 & -f_{12} & 0 & 0 \\ -f_{21} & 1 & 0 & 0 \end{pmatrix}. \quad (2.5)$$

It is possible to demonstrate that the real part of the quadratic form $(C^n)_i (M^n)_{ij} (C^n)_j$ in (2.4) is positive whenever the chemical potentials obey the inequalities

$$x < \min \left\{ y_1 y_2, \frac{y_1}{y_2}, \frac{y_2}{y_1}, \frac{1}{y_1 y_2} \right\} \quad (2.6)$$

a condition that it necessary for the index to be well defined in the first place ⁴ As a consequence it appears that the integral (2.3) is always dominated by the saddle point at $\rho_n = \chi_n = 0$ for all $n \geq 0$. It thus appears that, in perfect analogy with the situation for $\mathcal{N} = 4$ Yang Mills, the Witten index (2.3) never undergoes the phase transition into a black hole like phase.

As the saddle point contribution to the integral (2.3) vanishes, the first nonzero contribution to this integral is given by the inverse square root of the determinant

$$I^W = \prod_{n=1}^{\infty} \frac{1}{\sqrt{16 \det M^n}} \quad (2.7)$$

where the normalization in (2.7) is fixed by the requirement that I^W tends to unity when $x = 0$ (at which point only the vacuum contributes to the Witten index).

⁴In order that the index be well defined it is necessary that every $\Delta = 0$ letter contribute to the partition function with a weight less than unity, leading to (2.6).

The determinant is given by,

$$\begin{aligned}\det(M^n) &= \frac{1}{16}(1 - f_{12}(x^n, y_1^n, y_2^n)f_{21}(x^n, y_1^n, y_2^n))^2 \\ &= \frac{1}{16} \frac{\left(1 - \frac{x^n}{y_1^n}\right)^2 \left(1 - \frac{x^n}{y_2^n}\right)^2 (1 - x^n y_1^n)^2 (1 - x^n y_2^n)^2}{(1 - x^{2n})^4}\end{aligned}\quad (2.8)$$

so that,

$$I^W = \prod_n \frac{(1 - x^{2n})^2}{\left(1 - \frac{x^n}{y_1^n}\right) \left(1 - \frac{x^n}{y_2^n}\right) (1 - x^n y_1^n) (1 - x^n y_2^n)}.\quad (2.9)$$

As the large N unitary integrals in (2.3) never undergo a large N Gross-Witten-Wadia transition [15, 16], it follows that the Witten index I^W receives contributions only from states of finite energy (and charge) at finite values of the chemical potential. In particular, (2.9) is blind to states whose energy is of order N^a where a is any positive power.

In order to get a feel for (2.9) it is useful to set $y_1 = y_2 = 1$. If we define the Indicial entropy $S_{ind}(E)$ by the formula $I^w(x) = \int dE e^{S_{ind}(E)} x^E$ then it is easy to show that $S_{ind}(E) \approx \sqrt{2\pi} \sqrt{E}$ at high energies. This is the growth of states of a two dimensional massless gas; a similar growth in density of states was captured by the four dimensional index (see [7]). This growth is slower than the $E^{\frac{2}{3}}$ growth demonstrated by the index of the M2 brane and M5 brane world volume theories [3].

3. Gravity computation of the index

Gravitons⁵ in $AdS_4 \times S^7$ may be organized into representations of the $d = 3$ $\mathcal{N} = 8$ superconformal algebra. Working in conventions in which the $M2$ brane world volume scalar, fermion and supersymmetries respectively transform in the $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2})$, $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ and $(1, 0, 0, 0)$ representations of $SO(8)$, the highest weight states of the representations that occur in this decomposition each have $j = 0$, $\epsilon_0 = \frac{n}{2}$ and $SO(8)$ highest weight charges $(n/2, n/2, n/2, -n/2)$. See [3], Table 1 for more details.

Gravitons on $AdS_4 \times \frac{S^7}{Z_k}$ are those graviton states on $AdS_4 \times S^7$ whose charge under the generator $2h_4$ is 0 mod k .⁶ In the large k 't Hooft limit under study

⁵In this section we use the word ‘graviton’ for any field on AdS_4 obtained upon compactification from a field in the 11 dimensional gravity multiplet.

⁶As for $SO(6)$ we think of $SO(8)$ as the group of rotations in R^8 parameterized by x^i , $i = 1 \dots 8$. h_1, h_2, h_3, h_4 are the eigenvalues of the generators of rotations in the (12), (34), (56) and (78) planes respectively. We label representations of $SO(8)$ by their highest weights under (h_1, h_2, h_3, h_4) ; our positivity convention for weights is the obvious generalization of that for $SO(6)$. Note that the Z_k orbifolding described in [1] is, in our conventions, simply a rotation by the angle $4\pi/k$ in the (78) 2 plane. The subgroup of $SO(8)$ that commutes with this rotation is $SO(6) \times SO(2)$. The $SO(2)$

in this note, all projected in gravitons are simply neutral under the $U(1)$ charge h_4 . Consequently, the superconformal index over single gravitons in $AdS_4 \times \frac{S^7}{Z_k}$ is simply the projection of the same quantity in $AdS_4 \times S^7$ to the sector of zero h_4 charge.

The index

$$I^W = \text{Tr}[(-1)^F x^{\epsilon_0+j} y_1^{h_2} y_2^{h_3} y_3^{h_4}] \quad (3.1)$$

over single gravitons in $AdS_4 \times S^7$ was evaluated in [3] (see equation 2.17 in that paper). For the convenience of the reader we reproduce the formula here,

$$I_{AdS_4 \times S^7}^W(x, y_1, y_2, y_3) = \frac{\text{Numerator}}{\text{Denominator}}, \quad (3.2)$$

where,

$$\begin{aligned} \text{Numerator} = & -\sqrt{y_1}\sqrt{y_2}\sqrt{y_3}(y_2y_3y_1 + y_1 + y_2 + y_3)x^{7/2} \\ & + (y_2y_3y_1^2 + (y_3y_2^2 + y_3^2y_2 + y_2 + y_3)y_1 + y_2y_3)x^3 \\ & - (y_2y_3y_1^2 + (y_3y_2^2 + y_3^2y_2 + y_2 + y_3)y_1 + y_2y_3)x \\ & + \sqrt{y_1}\sqrt{y_2}\sqrt{y_3}(y_2y_3 + y_1(y_2 + y_3) + 1)\sqrt{x} \\ \text{Denominator} = & (1 - x^2)(\sqrt{x}\sqrt{y_1}\sqrt{y_2} - \sqrt{y_3})(\sqrt{x}\sqrt{y_1}\sqrt{y_3} - \sqrt{y_2}) \\ & (\sqrt{y_1} - \sqrt{x}\sqrt{y_2}\sqrt{y_3})(\sqrt{y_1}\sqrt{y_2}\sqrt{y_3} - \sqrt{x}) \end{aligned} \quad (3.3)$$

The index over single gravitons of zero h_4 charge is given by

$$\int \frac{d\theta}{2\pi i} I_{AdS_4 \times S^7}^W(x, y_1, y_2, e^{i\theta}) = \int_C \frac{dy_3}{2\pi y_3 i} I_{AdS_4 \times S^7}^W(x, y_1, y_2, y_3)$$

where the contour C surrounds the poles at $y_3 = 0$, $y_3 = xy_1y_2$ and $\frac{x}{y_1y_2}$.⁷

Performing this sum of residues we find that the index over $U(1)$ neutral gravitons on $AdS_4 \times \frac{S^7}{Z_k}$ is given by,

$$I_{\text{Single Particle}}^W = \frac{x}{y_1 - x} + \frac{1}{1 - xy_1} + \frac{x}{y_2 - x} + \frac{1}{1 - xy_2} - \frac{2}{1 - x^2}. \quad (3.4)$$

a result that is significantly simpler than (3.3).

We have also verified (3.4) more directly. As explained in [13, 1] the $U(1)$ neutral gravitons in $AdS_4 \times \frac{S^7}{Z_k}$ appear in a direct sum representations of the $\mathcal{N} = 6$ superconformal algebra labeled by the highest weight states with $\epsilon_0 = n$, $j = 0$ and $(h_1, h_2, h_3) = (n, n, 0)$ for $n = 1 \dots \infty$. It is not difficult to decompose every such

factor is simply rotations in the (78) 2 plane itself while the $SO(6)$ factor describes rotations among the remaining orthogonal 2 planes and is the R symmetry of the surviving $\mathcal{N} = 6$ supersymmetry algebra. Note that the supersymmetry of the parent theory decomposes into $6_0 + 1_2 + 1_{-2}$ under this decomposition, while the scalar decomposes into the $4_1 + \bar{4}_{-1}$.

⁷The index (3.3) is well defined only when $x < y_1^a y_2^b y_3^c$ where (a, b, c) run over the values of $(2h_1, 2h_2, 2h_3)$ for the antichiral spinor of $SO(6)$. It follows that the contour of our integral must exclude the poles at $y_3 = \frac{y_1}{xy_2}$ and $y_3 = \frac{y_2}{xy_1}$.

Range of n	Scaling dimension ϵ_0	$SO(3)$ weight (j)	$SO(6)$ highest weight (orthogonal basis)	$\epsilon_0 + j$	statistics
$n \geq 1$	n	0	$(n, n, 0)$	n	bosonic
$n \geq 1$	$n + \frac{1}{2}$	$\frac{1}{2}$	$(n, n, 1)$	$n + 1$	fermionic
$n \geq 1$	$n + \frac{1}{2}$	$\frac{1}{2}$	$(n, n, -1)$	$n + 1$	fermionic
$n \geq 1$	$n + \frac{1}{2}$	$\frac{1}{2}$	$(n, n - 1, 0)$	$n + 1$	fermionic
$n \geq 1$	$n + 1$	1	$(n, n, 0)$	$n + 2$	bosonic
$n \geq 2$	$n + 1$	1	$(n, n - 1, 1)$	$n + 2$	bosonic
$n \geq 2$	$n + 1$	1	$(n, n - 1, -1)$	$n + 2$	bosonic
$n \geq 2$	$n + \frac{3}{2}$	$\frac{3}{2}$	$(n, n - 1, 0)$	$n + 3$	fermionic

Table 3: The supersymmetric ($\Delta = 0$) graviton spectrum in $AdS_4 \times \frac{S^7}{Z_k}$. Here n is an integer greater than or equal to 1.

representation of the superconformal algebra into irreducible representations of the $d = 3$ conformal algebra (using, for instance, the techniques described in [3]). This decomposition could also be read off from the Table 1 in [13]. In Table 3 we list those conformal representations that have states with $\Delta = 0$. Only states with maximum values of h_1 and j in the representations listed in Table 3 have $\Delta = 0$ and contribute to the index. Consequently, the contribution of any of the representations listed below, to the index is simply given by

$$I_{\epsilon_0, j, h_2, h_3}^W = (-1)^j \frac{x^{\epsilon_0 + j}}{1 - x^2} \chi_{SO(4)}^{(h_2, h_3)}(y_1, y_2)$$

where ϵ_0 and j are respectively the dimension and $SO(3)$ charge of the highest weight state in the representation, the factor of $\frac{1}{1-x^2}$ is the contribution from the supersymmetric derivatives and $\chi_{SO(4)}^{(h_2, h_3)}(y_1, y_2)$ is the $SO(4)$ character with highest weights h_2, h_3 . Summing this quantity over all the representations listed in Table 3 for all $n \geq 1$ we recover (3.4).

In order to compute the index over multi gravitons we use the single particle index (3.4) and the formulas of Bose statistics to obtain ⁸

$$I^W = \prod_n \frac{(1 - x^{2n})^2}{\left(1 - \frac{x^n}{y_1^n}\right) \left(1 - \frac{x^n}{y_2^n}\right) (1 - x^n y_1^n) (1 - x^n y_2^n)}. \quad (3.5)$$

in perfect agreement with (2.9).

⁸According to the rules of Bose Statistics a single particle index of the form $\sum_n c_n x^n$ translates into a multi particle index $\prod_n (1 - x^n)^{-c_n}$ where c_n are integers that could be either negative or positive, and x schematically represents all chemical potentials.

4. Discussion

In this note we have computed the supersymmetric index of [3] for the ABJM theory in two different ways. We first performed a one loop field theory computation to evaluate this index in the free theory field at large N ; this calculation was performed at $\lambda = \frac{N}{k} = 0$. We then evaluated the same index over the Fock space of $U(1)$ neutral gravitons in $AdS_4 \times \frac{S^7}{\mathbb{Z}_k}$. This calculation was valid in the large N limit with λ fixed at a large value. The results of our two calculations match perfectly, providing a check of the ABJM conjecture.

It would be easy to generalize the gravitational calculation presented in this note to apply the large N limit with k held fixed. All one needs to do is to project (3.3) onto the sector with $2h_4 = 0 \bmod k$ (rather than simply to zero), before applying the formulas of Bose statistics. For instance when $k = 1$ this simply amounts to setting y_3 to unity in (3.3). It may be possible to reproduce the full finite k gravitational index from an (almost) free field theory calculation after summing over flux sectors on S^2 , as suggested by the discussion in [1]. It would be very interesting to try to carry this through.

Another direction that would be interesting to explore would be the determination of the full supersymmetric partition function (rather than the supersymmetric index) of the ABJM theory, with varying amounts of supersymmetry. For instance a formula for finite N partition function over the chiral ring (states that preserve 4 supercharges) has been proposed in [14] at $k = 1$. It would be interesting to verify this formula and to generalize it to other values of k . More ambitiously one could attempt to determine the full partition function (in contrast to the index computed in this paper) over all supersymmetric of in the ABJM theory; note however that this programme has not yet been completed even for $\mathcal{N} = 4$ Yang Mills (see [18] for a recent status report).

In this connection note that like $AdS_5 \times S^5$, the ABJM gravitational background presumably hosts supersymmetric black holes that preserve 2 supersymmetries (see [10, 11, 12] for relevant work). This is exactly the minimum amount of supersymmetry that a state needs to preserve to contribute to the index described in this paper. However the index computed in this paper sees no sign of these states. Indeed a naive estimate suggests that the entropy of these supersymmetric black holes scale like $N^2/\sqrt{\lambda}$ times functions of chemical potentials. As a result, the entropy of these black holes appears to be a function of λ at large λ , and so cannot be captured by any quantity like an index that is independent of λ . The smooth dependence of the entropy of a supersymmetric configuration on a continuous coupling constant appears non intuitive at first sight, and it would be interesting to understand how this comes about. Perhaps the states that make up the entropy of the black hole receive important contributions from the nontrivial flux sectors (these sectors, whose energy scales like N , could in principle contribute to the entropy of a black hole -

whose energy scales like N^2 , even in the 't Hooft limit).⁹

Turning to the spectrum of nonsupersymmetric states in this theory, it seems possible that the integrability of the spin chain spectrum of chiral operators could carry over to the ABJM theory. It would be interesting to study this possibility in more detail. Indeed, one of the exciting aspects of the ABJM proposal (in our opinion) are the prediction that the effective string that describes spin chain dynamics metamorphoses into a membrane at $\lambda = \mathcal{O}(N)$. It would be very interesting to attempt to get a concrete handle on this.

Finally, of course the ABJM duality permits the computation of all correlators (not just the spectrum) of all the chiral operators in the theory at strong coupling. A simple scaling estimate reveals that k point functions of these operators scale like $(\lambda^{\frac{1}{4}}/N)^{k-2}$ at strong coupling. In particular, three point functions are functions of λ and cannot enjoy the nonrenormalization properties of 3 point functions of chiral operators in $\mathcal{N} = 4$ Yang Mills theory in $d = 4$ [17].

Acknowledgements

We would especially like to thank S. Bhattacharyya and R. Loganayagam for several useful discussions during the course of this project. We would also like to thank M. Berkooz, N. Drukker, B. Ezhuthachan, M. Gaberdiel, R. Gopakumar, S. Kim, G. Mandal, S. Mukhi, C. Papageorgakis, S. Raju, S. Wadia, all the participants of Monsoon Workshop on String Theory and all the students in the TIFR theory room for useful discussions and comments on the manuscript. The work of SM was supported in part by a Swarnajayanti Fellowship. We would like to acknowledge our debt to the steady and generous support of the people of India to research in the basic sciences.

References

- [1] O. Aharony, O. Bergman, D. L. Jafferis and J. Maldacena, “ $N=6$ superconformal Chern-Simons-matter theories, M2-branes and their gravity duals”, [arXiv:0806.1218 [hep-th]]
- [2] M. Benna, I. Klebanov, T. Klose and M. Smedback, “Superconformal Chern-Simons Theories and AdS_4/CFT_3 Correspondence”, [[arXiv:0806.1519 [hep-th]].]
- [3] J. Bhattacharya, S. Bhattacharyya, S. Minwalla and S. Raju, “Indices for Superconformal Field Theories in 3,5 and 6 Dimensions”, *JHEP* **0802**, 064 (2008), [arXiv:0801.1435 [hep-th]]
- [4] B. Sundborg, “The Hagedorn transition, deconfinement and $N = 4$ SYM theory”, *Nucl. Phys. B* **573**, 349 (2000), [[arXiv:hep-th/9908001].]

⁹We thank R. Gopakumar for this suggestion.

- [5] O. Aharony, J. Marsano, S. Minwalla, K. Papadodimas and M. Van Raamsdonk, “*The Hagedorn / deconfinement phase transition in weakly coupled large N gauge theories*”, *Adv. Theor. Math. Phys.* **8**, 603 (2004) [[\[arXiv:hep-th/0310285\]](#) .]
- [6] O. Aharony, J. Marsano, S. Minwalla, K. Papadodimas and M. Van Raamsdonk, “*A first order deconfinement transition in large N Yang-Mills theory on a small S^3* ”, *Phys. Rev. D* **71**, 125018 (2005) [[\[arXiv:hep-th/0502149\]](#) .]
- [7] J. Kinney, J. M. Maldacena, S. Minwalla and S. Raju, “*An index for 4 dimensional super conformal theories*”, *Commun. Math. Phys.* **275**, 209 (2007) [[\[arXiv:hep-th/0510251\]](#) .]
- [8] Y. Nakayama, “*Index for orbifold quiver gauge theories*”, *Phys. Lett. B* **636**, 132 (2006) [[\[arXiv:hep-th/0512280\]](#) .]
- [9] D. Gaiotto and X. Yin, “*Notes on superconformal Chern-Simons-matter theories*”, *JHEP* **0708**, 056 (2007) [[\[arXiv:0704.3740 \[hep-th\]\]](#) .]
- [10] V. A. Kostelecky and M. J. Perry, “*Solitonic Black Holes in Gauged $N=2$ Supergravity*”, *Phys. Lett. B* **371**, 191 (1996) [[\[arXiv:hep-th/9512222\]](#) .]
- [11] Z. W. Chong, M. Cvetič, H. Lu and C. N. Pope, “*Charged rotating black holes in four-dimensional gauged and ungauged supergravities*”, *Nucl. Phys. B* **717**, 246 (2005) [[\[arXiv:hep-th/0411045\]](#) .]
- [12] M. Cvetič, G. W. Gibbons, H. Lu and C. N. Pope, “*Rotating black holes in gauged supergravities: Thermodynamics, supersymmetric limits, topological solitons and time machines*”, [[\[arXiv:hep-th/0504080\]](#) .]
- [13] B. E. W. Nilsson and C. N. Pope “*Hopf fibration of eleven-dimensional supergravity*”, *Class. Quantum Grav.* **1**(1984) 499-515.
- [14] S. Bhattacharyya and S. Minwalla, “*Supersymmetric states in $M5/M2$ CFTs*”, *JHEP* **0712**, 004 (2007) [[\[arXiv:hep-th/0702069\]](#) .]
- [15] D. J. Gross and E. Witten, “*Possible Third Order Phase Transition In The Large N Lattice Gauge Theory*”, *Phys. Rev. D* **21**, 446 (1980).
- [16] S. R. Wadia, “ *$N = \infty$ Phase Transition In A Class Of Exactly Soluble Model Lattice Gauge Theories*”, *Phys. Lett. B* **93**, 403 (1980).
- [17] S. Lee, S. Minwalla, M. Rangamani and N. Seiberg, “*Three-point functions of chiral operators in $D = 4$, $N = 4$ SYM at large N* ”, *Adv. Theor. Math. Phys.* **2**, 697 (1998) [[\[arXiv:hep-th/9806074\]](#) .]
- [18] L. Grant, P. A. Grassi, S. Kim and S. Minwalla, “*Comments on $1/16$ BPS Quantum States and Classical Configurations*”, *JHEP* **0805**, 049 (2008) [[\[arXiv:0803.4183 \[hep-th\]\]](#) .]